

DISEASES & CONDITIONS

Clavicle Fracture (Broken Collarbone)

A clavicle fracture is a break in the collarbone, one of the main bones in the shoulder girdle and chest. This type of [fracture](#) accounts for about 5% of all adult fractures.


Most clavicle fractures occur when a fall onto the shoulder or an outstretched arm puts enough pressure on the bone that it snaps or breaks. A broken collarbone can be very painful and can make it hard to move your arm.

- Many clavicle fractures can be treated by wearing a sling to keep the arm and shoulder from moving while the bone heals.
- With some clavicle fractures, however, the pieces of bone move far out of place when the injury occurs. For some of these more complicated fractures, surgery may be needed to realign the collarbone.

Anatomy

The clavicle is located between the sternum (ribcage) and the scapula (shoulder blade). It is the bone that connects the arm to the body.

The clavicle lies above several important nerves and blood vessels. Sometimes these vital structures are injured when a fracture occurs.



The clavicle is part of your shoulder and connects your arm to the rest of your body.

Reproduced and adapted from JF Sarwak, ed: Essentials of Musculoskeletal Care, ed. 4. Rosemont, IL, American Academy of Orthopaedic Surgeons, 2010.

Description


Clavicle fractures occur in people of all ages. Most fractures occur in the middle portion, or shaft, of the bone. Occasionally, the bone will break where it attaches at the ribcage or shoulder blade.

Clavicle fractures vary.

- The bone can crack just slightly or break into many pieces. This is called a comminuted fracture.
- The broken pieces of bone may line up straight or may be far out of place. This is called a displaced fracture.

This illustration shows a clavicle fracture close to where the bone attaches to the scapula (shoulder blade).

Reproduced and adapted from Nuber GW, Bowen MK: Acromioclavicular joint injuries and distal clavicle fractures. J Am Acad Orthop Surg 1997; 5(1): 11-18.



Cause

- Clavicle fractures are most often caused by a direct blow to the shoulder. This can happen if you fall directly onto the shoulder or have an accident (e.g., a motor vehicle collision or a sports injury).
- A fall onto an outstretched arm can also cause a clavicle fracture.
- In a baby, a clavicle fracture can occur during the passage through the birth canal.

Symptoms

A clavicle fracture can be very painful and may make it hard to move your arm. Other signs and symptoms of a fracture may include:

- Sagging of the shoulder downward and forward
- Inability to lift the arm because of pain
- A grinding sensation when you try to raise the arm
- A deformity or bump over the break
- Bruising, swelling, and/or tenderness over the collarbone

Doctor Examination

Physical Examination

Your doctor ask you how the injury occurred and about your symptoms. They will then carefully examine your shoulder.

In a clavicle fracture, there is usually an obvious deformity, or bump, at the fracture site. Gentle pressure over the break will bring about pain. Although it is rare for a bone fragment to break through the skin, the fragments may push on the skin. This is sometimes called "tenting" of the skin.



In a clavicle fracture, the broken ends of the bone may cause tenting of the skin over the fracture site.

Photo courtesy of Stuart J. Fischer, MD, FAAOS

Your doctor will also perform tests to make sure that no nerves or blood vessels were damaged when the fracture occurred.


Imaging Tests

X-rays. X-rays provide images of dense structures, such as bone. Your doctor will order an X-ray:

- To identify the location of the fracture
- To learn more about the severity of the break

The doctor may also order X-rays of your entire shoulder to check for additional injuries.

If other bones are broken, your doctor may order a computerized tomography (CT) scan to see the fractures in better detail.



X-ray shows a fracture in the middle of the clavicle. Note how far out of place (displaced) the broken ends of the bone are.

Image courtesy of Stuart J. Fischer, MD, FAAOS

Treatment

Nonsurgical Treatment

If the broken ends of the bones have not shifted much out of place, you may not need surgery. Many broken collarbones – especially in the middle portion, or shaft, of the collarbone – can heal without surgery.

Nonsurgical treatment may include:

- **Arm support.** A simple arm sling is usually used for comfort immediately after the break and while the injury heals.
- **Medication.** Pain medication, including acetaminophen or [nonsteroidal anti-inflammatory medications \(NSAIDs\)](#), can help relieve pain as the fracture heals.
- **Motion and exercises.** Although there will be some pain, it is important to maintain arm motion to prevent shoulder and elbow stiffness. Often, patients will begin doing exercises for hand, wrist, and elbow motion immediately after the injury. Your doctor may order formal physical therapy for you.

After a clavicle fracture, once the bone begins to heal, your pain will decrease and your doctor may start you on gentle shoulder exercises. These exercises will help prevent stiffness and weakness. You will start more strenuous exercises gradually, once the fracture is completely healed.

Follow-up care. You will need to see your doctor regularly until your fracture heals. During these visits, the doctor will take X-rays to make sure the bone is healing in a good position. After the bone has healed, you will be able to gradually return to your normal activities.

Complications. In some cases, a clavicle fracture can move out of place before it heals. It is important to follow up with your doctor as scheduled to make sure the bone stays in position.

If the fracture does not heal, it is called a [nonunion](#).

- In some cases of nonunion, the patient has very little pain and good use of their arm, so no further treatment is required.
- In other cases, nonunion can result in pain and arm dysfunction and may require an operation for repair.

If the fracture fragments do move out of place and the bones heal in that position, it is called a malunion. Surgical treatment for this is very rare. The decision to perform surgery is based on how far out of place the bones are and how much the displacement affects your arm movement.

A large bump over the fracture site may develop as the fracture heals. This usually gets smaller over time, but a small bump often remains permanently.

Surgical Treatment

If the broken ends of the bones are very displaced (out of normal alignment), your doctor may recommend surgery.

Surgery typically involves putting the broken pieces of bone back into their proper position and preventing them from moving out of place until they are healed. This can improve shoulder strength when you have recovered.


Open reduction and internal fixation. This is the procedure most often used to treat clavicle fractures. During the procedure, the surgeon makes an incision (a cut) over the bone and first reduces (repositions) the bone fragments into their normal alignment. The pieces of bone are then held in place with special metal hardware.

Common methods of internal fixation include:

- **Plates and screws.** After being repositioned into their normal alignment, the bone fragments are held in place with special screws and metal plates attached to the outer surface of the bone.

After surgery, you may notice a small patch of numb skin below the incision. This numbness will become less noticeable with time. Because the clavicle lies directly under the skin, you may be able to feel the plate through your skin.

Plates and screws are not usually removed after the bone has healed, unless they are causing discomfort. Problems with the hardware are not common, but some patients find that seatbelts and backpacks can irritate the collarbone area. If this happens, the hardware can be removed after the fracture has healed.




(Left) X-ray shows a displaced clavicle fracture (arrow). **(Right)** The pieces of bone have been realigned and held in place with plates and screws.

Images courtesy of Stuart J. Fischer, MD, FAAOS

- **Pins or screws.** Pins or screws can also be used to hold the fracture in good position after the bone ends have been put back in place. The incisions for pin or screw placement are usually smaller than those used for plates.

Pins or screws often irritate the skin where they have been inserted and are usually removed once the fracture has healed.



(Left) X-ray shows a severely displaced clavicle fracture (arrow). **(Right)** Here, a single screw has been used to repair the fracture.

Reproduced from Eichinger JK, Balog TP, Grassbaugh JA: Intramedullary fixation of clavicle fractures: anatomy, indications, advantages, and disadvantages. J Am Acad Orthop Surg 2016; 24(7): 455-464.

Pain management. After surgery, you will feel some pain. This is a natural part of the healing process. Ice and non-prescription pain medications are often enough to relieve pain.

In some cases, your doctor may suggest a prescription-strength medication, such as an opioid, for a few days.

Be aware that although opioids help relieve pain after surgery, their use has risks and complications. These medications can be addictive and potentially dangerous. It is therefore important to use opioids only as directed by your doctor, to use as little as possible for as short a time as possible, and to stop taking them as soon as your pain starts to improve. Tell your doctor if your pain has not begun to improve within a few days after surgery.

Rehabilitation. Specific exercises will help restore movement and strengthen your shoulder. Your doctor may provide you with a home therapy plan that you do on your own or suggest that you work with a therapist.

Therapy programs typically start with gentle motion exercises. Your doctor will gradually add strengthening exercises to your program as your fracture heals.

Although it is a slow process, following your therapy plan is an important factor in returning to all the activities you enjoy.

Complications. There are risks associated with any type of surgery. These include:

- [Infection](#)
- Bleeding
- Problems with wound healing
- Pain
- [Blood clots](#)
- Damage to blood vessels or nerves
- Reaction to anesthesia

Risks that are specific to surgery for clavicle fractures include:

- Difficulty with bone healing
- Lung injury
- Numbness below the clavicle

- Hardware irritation

Patients who smoke or use tobacco products, have diabetes, or are elderly have a higher risk of complications both during and after surgery. They are also more likely to have problems with wound and bone healing.

Learn more: [Smoking and Musculoskeletal Health](#)

Learn more: [Surgery and Smoking](#)

Learn more: [Impact of Chronic Illness on Bone Health](#)

Before your surgery, your doctor will discuss each of the risks with you and will take specific measures to avoid complications.

Outcome

Whether or not your treatment involves surgery, it can take several months for your collarbone to heal. Healing may take longer in diabetics or in people who smoke or use tobacco products.

Most people return to their regular activities within 3 months of their injury. Your doctor will tell you when your injury is stable enough to do so. Returning to regular activities or lifting with your arm before your doctor advises may cause the fracture fragments to move or the hardware to break. This may require you to start your treatment from the beginning.

Once your fracture has completely healed, you can safely return to sports and other physical activities.

To assist doctors in the management of clavicle fractures, the American Academy of Orthopaedic Surgeons has conducted research to provide some useful guidelines. These are recommendations only and may not apply to every case. For more information: [Plain Language Summary - Clinical Practice Guideline - The Treatment of Clavicle Fractures - AAOS](#)

Last Reviewed

September 2025

Contributed and/or Updated by

[Gillian LS Soles, MD, FAAOS](#)

Peer-Reviewed by

[Julie E. Adams, MD, FAAOS](#)

AAOS does not endorse any treatments, procedures, products, or physicians referenced herein. This information is provided as an educational service and is not intended to serve as medical advice. Anyone seeking specific orthopaedic advice or assistance should consult his or her orthopaedic surgeon, or locate one in your area through the AAOS [Find an Orthopaedist](#) program on this website.

